

MAPK3

Reactivity:Human Mouse

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:1000

Calculated MW:44kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human MAPK3

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

ioqr

Synonym:

MAPK3;ERK1;HS44KDAP;HUMKER1A;MGC20180;P44ERK1;P44MAPK;PRKM3 ;ERK1/2

Catalog #:A0025

Antibody Type:

Monoclonal Antibody

Species:Mouse

Gene ID:5595

Isotype:IgG

Swiss Prot:P27361

Purity:Affinity purification

For research use only.

Background:

Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (MAPK3/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli including mitogens, growth factors, and cytokines (1-3) and is an important target in the diagnosis and treatment of cancer (4). Upon stimulation, a sequential three-part protein kinase cascade is initiated, consisting of a MAP kinase kinase kinase (MAPKKK or MAP3K), a MAP kinase kinase (MAPKK or MAP2K), and a MAP kinase (MAPK). Multiple p44/42 MAP3Ks have been identified, including members of the Raf family as well as Mos and Tpl2/Cot. MEK1 and MEK2 are the primary MAPKKs in this pathway (5,6). MEK1 and MEK2 activate p44 and p42 through phosphorylation of activation loop residues Thr202/Tyr204 and Thr185/Tyr187, respectively. Several downstream targets of p44/42 have been identified, including p90RSK (7) and the transcription factor Elk-1 (8,9). p44/42 are negatively regulated by a family of dual-specificity (Thr/Tyr) MAPK phosphatases, known as DUSPs or MKPs (10), along with MEK inhibitors such as U0126 and PD98059.

To place an order, please [Click HERE](#).