

## CYP1A2

**Reactivity:**Mouse Rat

**Tested applications:**WB

**Recommended Dilution:**WB 1:500 - 1:2000

**Calculated MW:**58kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant Protein of human CYP1A2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CP12; P3-450; P450(PA);

**Catalog #:**A0062

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**1544

**Isotype:**IgG

**Swiss Prot:**P05177

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. The protein encoded by this gene localizes to the endoplasmic reticulum and its expression is induced by some polycyclic aromatic hydrocarbons (PAHs), some of which are found in cigarette smoke. The enzyme's endogenous substrate is unknown; however, it is able to metabolize some PAHs to carcinogenic intermediates. Other xenobiotic substrates for this enzyme include caffeine, aflatoxin B1, and acetaminophen. The transcript from this gene contains four Alu sequences flanked by direct repeats in the 3' untranslated region.

**To place an order, please [Click HERE](#).**