www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

BCR



Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW:130/160kDa Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human BCR

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

ALL;BCR1;CML;D22S11;D22S662;PHL;Breakpoint cluster region protein antibody, Renal carcinoma antigen NY-REN-26 antibody;

Background:

The breakpoint cluster region protein (Bcr) is best know to be involved in genomic translocation with fusion partner Abl (Cbr-Abl) causing chronic myelogenous leukemia (CML). This 160 kDa protein contains a serine/threonine kinase domain, an SH2 binding domain, a GTP/GDP exchange domain and a C-term domain which functions as a GTPase activating protein for p21rac and CDC42. Additionally, Bcr is involved in signal transduction and can down regulate Ras mediated cell signaling.

To place an order, please Click HERE.

Catalog #:A0068

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:613

Isotype:IgG

Swiss Prot:P11274

Purity: Affinity purification

For research use only.





