

BCL2A1

Reactivity:Human Mouse

Tested applications:WB IHC FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 FC1:20 - 1:50

Calculated MW:20kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human BCL2A1

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

b

Synonym:

GRS; BFL1; ACC-1; ACC-2; HBPA1; BCL2L5;

Catalog #:A0134

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:597

Isotype:IgG

Swiss Prot:Q16548

Purity:Affinity purification

For research use only.

Background:

This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and pro-apoptotic regulators that are involved in a wide variety of cellular activities such as embryonic development, homeostasis and tumorigenesis. The protein encoded by this gene is able to reduce the release of pro-apoptotic cytochrome c from mitochondria and block caspase activation. This gene is a direct transcription target of NF-kappa B in response to inflammatory mediators, and is up-regulated by different extracellular signals, such as granulocyte-macrophage colony-stimulating factor (GM-CSF), CD40, phorbol ester and inflammatory cytokine TNF and IL-1, which suggests a cytoprotective function that is essential for lymphocyte activation as well as cell survival. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).