

## SPHK1

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**43kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human SPHK1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

ab

**Synonym:**

PHK1; SPHK

**Catalog #:**A0139

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**8877

**Isotype:**IgG

**Swiss Prot:**Q9NYA1

**Purity:**Affinity purification

For research use only.

**Background:**

Sphingosine kinases (SPHKs) catalyze the phosphorylation of sphingosine to form sphingosine-1-phosphate (S1P), a lipid mediator with both intra- and extracellular functions. Together with other sphingolipid metabolizing enzymes, SPHKs regulate the balance of the lipid mediators, ceramide, sphingosine, and S1P (1-4). Two distinct SPHK isoforms, SPHK1 and SPHK2, have been cloned and characterized (5,6). SPHK1 and SPHK2 are highly conserved and diversely expressed (7,8). The SPHKs are activated by G-protein-coupled receptors, receptor tyrosine kinases, immunoglobulin receptors, cytokines, and other stimuli (9-12). The molecular mechanisms by which SPHK1 and SPHK2 are specifically regulated are complex and only partially understood.

*To place an order, please [Click HERE](#).*