www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

CRP

Reactivity: Human Mouse

Tested applications: WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

Calculated MW:25kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human CRP

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

CRP;MGC149895;MGC88244;PTX1;

Background:

Human c-Reactive protein (CRP) was discovered in serum of patients as a nonantibody precipitin of the C-polysaccharide from pneumococci(1). During acute infection, the amounts of circulating CPR increases 100-1000 fold (2). Serum CRP values are routinely measured to detect or monitor many human diseases and minor elevations of C-reactive protein are predictive of cardiovascular events in patients with coronary heart disease (3). CRP promotes host defense, is an efficient activator of the classical C cascade, and shown to be opsonic (4). The opsonic and C activation activities are responsible for the nonspecific host defense activities providing protection in the preimmune stage of acute infections. Additionally, IL-1 and IL-6 are known to stimulate CRP biosynthesis (5).

To place an order, please Click HERE.

Species: Rabbit

Gene ID:1401

Isotype:IgG

Swiss Prot:P02741

Purity: Affinity purification

For research use only.





