CTNNB1

Reactivity: Human Mouse Rat

Tested applications: WB IHC ICC IP

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IP 1:20 - 1:50

Calculated MW:92kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human CTNNB1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

at

Synonym:

CTNNB1; CTNNB; DKFZp686D02253; FLJ25606; FLJ37923; Catenin beta-1; Beta-catenin ;-Catenin;

Background:

-catenin is a key downstream effector in the Wnt signaling pathway (1). It is implicated in two major biological processes in vertebrates: early embryonic development (2) and tumorigenesis (3). CK1 phosphorylates -catenin at Ser45. This phosphorylation event primes -catenin for subsequent phosphorylation by GSK-3 (4-6). GSK-3 destabilizes -catenin by phosphorylating it at Ser33, Ser37, and Thr41 (7). Mutations at these sites result in the stabilization of -catenin protein levels and have been found in many tumor cell lines (8).

To place an order, please Click HERE.

Gene ID:1499

Isotype:IgG

Swiss Prot:P35222

Purity: Affinity purification

For research use only.







