

## AKT2

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**56kDa

**Observed MW:**Refer to Figures

**Immunogen:**

A synthetic peptide of human AKT2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

opr

**Synonym:**

AKT2;PKBB;PKBBETA;PRKBB;RAC-BETA ;

**Catalog #:**A0336

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**208

**Isotype:**IgG

**Swiss Prot:**P31751

**Purity:**Affinity purification

For research use only.

**Background:**

Akt, also referred to as PKB or Rac, plays a critical role in controlling survival and apoptosis (1-3). This protein kinase is activated by insulin and various growth and survival factors to function in a wortmannin-sensitive pathway involving PI3 kinase (2,3). Akt is activated by phospholipid binding and activation loop phosphorylation at Thr308 by PDK1 (4) and by phosphorylation within the carboxy terminus at Ser473. The previously elusive PDK2 responsible for phosphorylation of Akt at Ser473 has been identified as mammalian target of rapamycin (mTOR) in a rapamycin-insensitive complex with rictor and Sin1 (5,6). Akt promotes cell survival by inhibiting apoptosis through phosphorylation and inactivation of several targets, including Bad (7), forkhead transcription factors (8), c-Raf (9), and caspase-9. PTEN phosphatase is a major negative regulator of the PI3 kinase/Akt signaling pathway (10). LY294002 is a specific PI3 kinase inhibitor (11). Another essential Akt function is the regulation of glycogen synthesis through phosphorylation and inactivation of GSK-3 and (12,13). Akt may also play a role in insulin stimulation of glucose transport (12). In addition to its role in survival and glycogen synthesis, Akt is involved in cell cycle regulation by preventing GSK-3-mediated phosphorylation and degradation of cyclin D1 (14) and by negatively regulating the cyclin dependent kinase inhibitors p27 Kip (15) and p21 Waf1/CIP1 (16). Akt also plays a critical role in cell growth by directly phosphorylating mTOR in a rapamycin-sensitive complex containing raptor (17). More importantly, Akt phosphorylates and inactivates tuberin (TSC2), an inhibitor of mTOR within the mTOR-raptor complex (18,19).

**To place an order, please [Click HERE](#).**