

GCK

Reactivity:Human Mouse

Tested applications:WB IHC IF

Recommended Dilution:WB 1:200 - 1:500 IHC 1:50 - 1:100 IF 1:20 - 1:50

Calculated MW:52kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human GCK

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

GCK;GK;GLK;HHF3;HK4;HKIV;HXKP;MODY2 ;

Catalog #:A0430

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:2645

Isotype:IgG

Swiss Prot:P35557

Purity:Affinity purification

For research use only.

Background:

Germinal center kinase (GCK) is the founding member of the GCK family, a group of serine/threonine kinases homologous to the yeast Ste20 kinase. GCK is activated by TNF and associated with TRAF2 (TNF receptor-associated factor 2) and MEKK1, thereby activating the SAPK/JNK pathway (1,2). GCK does not significantly phosphorylate MEKK1, instead it enhances MEKK1 oligomerization and autophosphorylation (3). GCK binds and activates MLK3, indicating that GCK may have other effectors (3).

To place an order, please [Click HERE](#).