

TCL1A

Reactivity:Human Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

Calculated MW:13kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human TCL1A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

TCL1A;TCL1 ;

Catalog #:A0629

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8115

Isotype:IgG

Swiss Prot:P56279

Purity:Affinity purification

For research use only.

Background:

TCL1 (T cell leukemia 1), MTCP1 and TCL1b belong to the TCL1 proto-oncogene family, and their products are involved in Akt activation during embryonic development, T cell leukemias, prolymphocytic leukemias and B cell lymphomas (1-3). The Akt association domain of TCL1 binds with the PH domain of Akt. The formation of an oligomeric TCL-Akt complex is required for TCL1 coactivator function and results in phosphorylation and activation of Akt . Furthermore, functional analysis indicates that the interaction between TCL1 and Akt promotes translocation of Akt to the nucleus (4-6). These findings are supported by the crystal structure of TCL1, which suggests that TCL1 may participate in molecular transport (7).

To place an order, please [Click HERE](#).