

TNFRSF6B

Reactivity:Human Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:33kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human TNFRSF6B

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bp

Synonym:

TNFRSF6B;DCR3; TR6

Catalog #:A0649

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8771

Isotype:IgG

Swiss Prot:O95407

Purity:Affinity purification

For research use only.

Background:

The tumor necrosis factor receptor family, which includes TNF-R1, Fas, DR3, DR4, DR5, and DR6, plays an important role in the regulation of apoptosis in various physiological systems (1,2). The receptors are activated by a family of cytokines that include TNF, FasL, and TRAIL. They are characterized by a highly conserved extracellular region containing cysteine-rich repeats and a conserved intracellular region of about 80 amino acids termed the death domain (DD). The DD is important for transducing the death signal by recruiting other DD containing adaptor proteins (FADD, TRADD, RIP) to the death-inducing signaling complex (DISC), resulting in activation of caspases. Death receptor signaling is also controlled by a family of decoy receptors (DcR1, DcR2 and DcR3) which lack a cytoplasmic DD and inhibit death receptor-mediated apoptosis by competing for ligand (3-5). Expression of decoy receptors provide a mechanism for certain types of cancer to regulate apoptosis and can contribute to chemosensitivity (6-8).

To place an order, please [Click HERE](#).