www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

RIPK3

Reactivity: Human Mouse

Tested applications: WB IP

Recommended Dilution:WB 1:200 - 1:2000 IP 1:20 - 1:50

Calculated MW:57kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human RIPK3

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

bp

Synonym:

RIPK3;RIP3;

Background:

The product of this gene is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is predominantly localized to the cytoplasm, and can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor.

To place an order, please Click HERE.

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:11035

Isotype:IgG

Swiss Prot:Q9Y572

Purity: Affinity purification

For research use only.





