

ATG7

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:78kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human ATG7

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

ch

Synonym:

ATG7;APG7-LIKE;APG7L;DKFZp434N0735;GSA7 ;

Catalog #:A0691

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:10533

Isotype:IgG

Swiss Prot:O95352

Purity:Affinity purification

For research use only.

Background:

Autophagy is a catabolic process for the autophagosomic-lysosomal degradation of bulk cytoplasmic contents (1,2). Autophagy is generally activated by conditions of nutrient deprivation but has also been associated with a number of physiological processes including development, differentiation, neurodegeneration, infection, and cancer (3). The molecular machinery of autophagy was largely discovered in yeast and referred to as autophagy-related (Atg) genes. Formation of the autophagosome involves a ubiquitin-like conjugation system in which Atg12 is covalently bound to Atg5 and targeted to autophagosome vesicles (4-6). This conjugation reaction is mediated by the ubiquitin E1-like enzyme Atg7 and the E2-like enzyme Atg10 (7,8).

To place an order, please [Click HERE](#).