

ANGPT2

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:57kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human ANGPT2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

a

Synonym:

ANG-2; ANG2; AGPT2; Angiopoietin-2;

Catalog #:A0698

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:285

Isotype:IgG

Swiss Prot:O15123

Purity:Affinity purification

For research use only.

Background:

Angiopoietins are a family of Tie receptor ligands. There are four angiopoietins discovered so far: angiopoietins 1, 2, 3 and 4 (Ang1, 2, 3, and 4) (1-3). Ang1 binds to the Tie-2 receptor and leads to its autophosphorylation and subsequent activation of downstream signaling pathways. It plays an important role in blood vessel formation, maturation and subsequent stabilization (1,4,5). Ang2 is an endothelium-specific growth factor that functions as an antagonist to Ang1, promotes vascular associated proinflammatory function, destabilizes quiescent endothelium, leads to vascular leakage and vascular destabilization and remodeling (2,6,7). Ang2 is selectively expressed in many tumor tissues where, combined with other growth factors such as VEGF, it can promote vascular remodeling, angiogenesis and inflammation (7-9).

To place an order, please [Click HERE](#).