

CHEK1

Reactivity:Human Mouse

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:1000 IHC 1:50 - 1:100

Calculated MW:54kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human CHEK1

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

e

Synonym:

CHK1

Catalog #:A0700

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:1111

Isotype:IgG

Swiss Prot:O14757

Purity:Affinity purification

For research use only.

Background:

In response to DNA damage, mammalian cells prevent cell cycle progression through the control of critical cell cycle regulators. CHK1 (synonym: CHEK1), a homolog of the *Schizosaccharomyces pombe* Chk1 protein kinase, is required for the DNA damage checkpoint. Human Chk1 protein is modified in response to DNA damage. In vitro Chk1 binds to and phosphorylate the dual-specificity protein phosphatases Cdc25A, Cdc25B, and Cdc25C, which control cell cycle transitions by dephosphorylating cyclin-dependent kinases. CHK1 can be autophosphorylated (PMID:22941630) and ubiquitinated (PMID:19276361). It has 3 isoforms produced by alternative splicing with the molecular weight of 54 kDa, 44 kDa and 50 kDa. This antibody is specific to CHK1.

To place an order, please [Click HERE](#).