

ATF2

Reactivity:Human Mouse

Tested applications:WB

Recommended Dilution:WB 1:200 - 1:500

Calculated MW:23kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human ATF2

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fk

Synonym:

ATF2;CRE-BP1;CREB2;HB16;MGC111558;TREB7

Catalog #:A0756

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:1386

Isotype:IgG

Swiss Prot:P15336

Purity:Affinity purification

For research use only.

Background:

The transcription factor ATF-2 (also called CRE-BP1) binds to both AP-1 and CRE DNA response elements and is a member of the ATF/CREB family of leucine zipper proteins (1). ATF-2 interacts with a variety of viral oncoproteins and cellular tumor suppressors and is a target of the SAPK/JNK and p38 MAP kinase signaling pathways (2-4). Various forms of cellular stress, including genotoxic agents, inflammatory cytokines, and UV irradiation, stimulate the transcriptional activity of ATF-2. Cellular stress activates ATF-2 by phosphorylation of Thr69 and Thr71 (2-4). Both SAPK and p38 MAPK have been shown to phosphorylate ATF-2 at these sites in vitro and in cells transfected with ATF-2. Mutations of these sites result in the loss of stress-induced transcription by ATF-2 (2-4). In addition, mutations at these sites reduce the ability of E1A and Rb to stimulate gene expression via ATF-2 (2).

To place an order, please [Click HERE](#).