

VDAC1

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:31kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human VDAC1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

ghi

Synonym:

VDAC1;MGC111064;PORIN;PORIN-31-HL ;

Catalog #:A0810

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7416

Isotype:IgG

Swiss Prot:P21796

Purity:Affinity purification

For research use only.

Background:

Voltage-dependent anion channel (VDAC), ubiquitously expressed and located in the outer mitochondrial membrane, is generally thought to be the primary means by which metabolites diffuse in and out of the mitochondria (1). In addition, this channel plays a role in apoptotic signaling. The change in mitochondrial permeability characteristic of apoptosis is mediated by Bcl-2 family proteins, which bind to VDAC, altering the channel kinetics (2). Homodimerization of VDAC may be a mechanism for changing mitochondrial permeability and supporting release of cytochrome c (3). In mammalian cells, there are three VDAC isoforms, VDAC1, which is the most widely expressed isoform, as well as VDAC2 and VDAC3 (4,5).

To place an order, please [Click HERE](#).