

CTTN

Reactivity:Human

Tested applications:WB IHC

Recommended Dilution:WB 1:1000 - 1:2000 IHC 1:50 - 1:100

Calculated MW:57kD

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human CTTN

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

EMS1;FLJ34459;

Catalog #:A0841

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:2017

Isotype:IgG

Swiss Prot:Q14247

Purity:Affinity purification

For research use only.

Background:

This gene is overexpressed in breast cancer and squamous cell carcinomas of the head and neck.

The encoded protein is localized in the cytoplasm and in areas of the cell-substratum contacts.

This gene has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells.

During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of this gene contributes to tumor cell invasion and metastasis.

Three splice variants that encode different isoforms have been identified for this gene.

To place an order, please [Click HERE](#).