

IKBKG

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:100 IF 1:50 - 1:200

Calculated MW:48kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human IKBKG

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bps

Synonym:

IKBKG;AMCBX1;FIP-3;FIP3;Fip3p;IKK-gamma;IP;IP1;IP2;IPD2;NEMO ; IKK;

Catalog #:A0917

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8517

Isotype:IgG

Swiss Prot:Q9Y6K9

Purity:Affinity purification

For research use only.

Background:

The NF- κ B/Rel transcription factors are present in the cytosol in an inactive state, complexed with the inhibitory IB proteins (1-3). Most agents that activate NF- κ B do so through a common pathway based on phosphorylation-induced, proteasome-mediated degradation of IB (3-7). The key regulatory step in this pathway involves activation of a high molecular weight IB kinase (IKK) complex whose catalysis is generally carried out by three tightly associated IKK subunits. IKK and IKK serve as the catalytic subunits of the kinase and IKK serves as the regulatory subunit (8,9). Activation of IKK depends upon phosphorylation of Ser177 and Ser181 in the activation loop of IKK (Ser176 and Ser180 in IKK), which causes conformational changes resulting in kinase activation (10-13).

To place an order, please [Click HERE](#).