

## PPIA

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**18kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human PPIA

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

PPIA;CYPA;CYPH;MGC117158;MGC12404;MGC23397 ;

**Catalog #:**A0993

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5478

**Isotype:**IgG

**Swiss Prot:**P62937

**Purity:**Affinity purification

For research use only.

**Background:**

Cyclophilins are a highly conserved family of peptidylprolyl cis-trans-isomerases (PPIA) that are targets of the immunosuppressant drug cyclosporin A (CsA) (1,2). The complex of cyclophilin and CsA can bind to and inhibit calcineurin which leads to inhibition of the transcription factor NFAT and decreased production of cytokines (3,4). As isomerases, cyclophilins have been proposed to aid in protein folding. Cyclophilin A can bind to the p55 Gag protein of HIV and appears necessary for HIV infection (5,6). There is also some evidence that cyclophilins have nuclease activity and play a role in apoptosis (7).

**To place an order, please [Click HERE](#).**