

ID2

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:18kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human ID2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

k

Synonym:

ID2;GIG8;ID2A;ID2H;MGC26389;bHLHb26 ;

Catalog #:A0996

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3398

Isotype:IgG

Swiss Prot:Q02363

Purity:Affinity purification

For research use only.

Background:

Inhibitor of DNA-binding-2 (Id2) is a member of the Id proteins which belong to the helix-loop-helix (HLH) protein family. The Id protein functions by binding to specific transcription factors and preventing their dimerization and DNA binding (1-3). Id2 interacts with a wide variety of transcription factors including E proteins (5), TCS (4), Pax (6) and the tumor suppressor Rb (1). Id2 has been shown to be important in regulating cellular differentiation, proliferation, development and tumorigenesis (7-9). In tumor cells, increased levels of Id2 functionally inactivate Rb, leading to cellular transformation and cancer (10,11). Id2 is therefore a promising therapeutic target for treatment of cancer (12).

To place an order, please [Click HERE](#).