www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## NCK1

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:43kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human NCK1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

NCK1;MGC12668;NCK;NCKalpha;

Background:

NCK1 (also known as NCK or NCK) is a broadly expressed oncogenic adapter protein consisting of three SH3 domains and one SH2 domain (1-3). NCK1 becomes phosphorylated upon activation of variety of cell surface receptors and is involved in actin cytoskeletal organization induced by many stimuli (4-6). NCK2 (also known as NCK), a homolog of NCK1, has an overlapping expression pattern and redundant functions with NCK1 (7).

To place an order, please Click HERE.

Polyclonal Antibody

Species:Rabbit

Gene ID:4690

Isotype:IgG

Swiss Prot:P16333

Purity: Affinity purification

For research use only.



