

YAP1

Reactivity: Human Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW: 75kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human YAP1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fn

Synonym:

YAP; YKI; YAP2; YAP65;

Catalog #: A1001

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 10413

Isotype: IgG

Swiss Prot: P46937

Purity: Affinity purification

For research use only.

Background:

The transcriptional coactivator Yes-associated protein (YAP) has been shown to interact with and to enhance p73-dependent apoptosis in response to DNA damage. It has been shown that YAP requires the promyelocytic leukemia gene (PML) and nuclear body localization to coactivate p73. YAP imparts selectivity to p73 by promoting the activation of a subset of p53 and/or p73 target promoters (1). Akt promotes YAP localization to the cytoplasm, resulting in loss from the nucleus where it functions as a coactivator of transcription factors including p73. p73-mediated induction of Bax expression following DNA damage requires YAP function and is attenuated by Akt phosphorylation of YAP. YAP overexpression increases, while YAP depletion decreases, p73-mediated apoptosis following DNA damage, in an Akt inhibitable manner (2)

To place an order, please [Click HERE](#).