

PAX8

Reactivity:Human

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000

Calculated MW:48kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human PAX8

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

k

Synonym:

Paired box 8; Paired box protein Pax 8; PAX8;

Catalog #:A1009

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7849

Isotype:IgG

Swiss Prot:Q06710

Purity:Affinity purification

For research use only.

Background:

PAX8 is a member of the paired box (PAX) family of transcription factors, typically containing a paired box domain and a paired-type homeodomain. It is expressed during organogenesis of the thyroid gland, kidney and Mullerian system. It is thought to regulate the expression of Wilms tumor suppressor (WT1) gene and mutations in PAX8 have been associated with Wilms tumor cells, thyroid and ovarian carcinomas. PAX8 is a useful marker in distinguishing ovarian carcinomas from mammary carcinomas (PMID: 18724243). PAX8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX8 has 5 isoforms with calculated MW 31kd, 34kd, 41kd, 43kd and 48kd. And there is a 60kd band corresponding to the electrophoretic mobility of PAX8 (PMID:15650356). Proteintechs 10336-1-AP antibody has cross-reactivities with the other PAX family members.

To place an order, please [Click HERE](#).