

## ASIC3

**Reactivity:**Mouse Rat

**Tested applications:**WB

**Recommended Dilution:**WB 1:500 - 1:2000

**Calculated MW:**59kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human ASIC3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

ACCN3; TNaC1; DRASIC; SLNAC1;

**Catalog #:**A10268

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**9311

**Isotype:**IgG

**Swiss Prot:**Q9UHC3

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, two hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene is an acid sensor and may play an important role in the detection of lasting pH changes. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 2 has been observed as proton-gated channels sensitive to gadolinium. Alternatively spliced transcript variants have been described.

**To place an order, please [Click HERE](#).**