

## SGCG

**Reactivity:** Human

**Tested applications:** WB

**Recommended Dilution:** WB 1:1000 - 1:2000

**Calculated MW:** 32kDa

**Observed MW:** Refer to figures

**Immunogen:**

Recombinant protein of human SGCG

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

A4; MAM; DMDA; SCG3; 35DAG; DAGA4; DMDA1; LGMD2C; SCARMD2; gamma-SG;

**Catalog #:** A10540

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 6445

**Isotype:** IgG

**Swiss Prot:** Q13326

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes gamma-sarcoglycan, one of several sarcolemmal transmembrane glycoproteins that interact with dystrophin. The dystrophin-glycoprotein complex (DGC) spans the sarcolemma and is comprised of dystrophin, syntrophin, alpha- and beta-dystroglycans and sarcoglycans. The DGC provides a structural link between the subsarcolemmal cytoskeleton and the extracellular matrix of muscle cells. Defects in the encoded protein can lead to early onset autosomal recessive muscular dystrophy, in particular limb-girdle muscular dystrophy, type 2C (LGMD2C).

**To place an order, please [Click HERE](#).**