www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

SCN2A

Reactivity: Human Mouse Rat

Tested applications:WB

Recommended Dilution: WB 1:1000 - 1:2000

Calculated MW:228kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human SCN2A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

HBA; NAC2; BFIC3; BFIS3; BFNIS; HBSCI; EIEE11; HBSCII; Nav1.2; SCN2A1; SCN2A2;

Na(v)1.2;

Background:

Voltage-gated sodium channels are transmembrane glycoprotein complexes composed of a large alpha subunit with 24 transmembrane domains and one or more regulatory beta subunits. They are responsible for the generation and propagation of action potentials in neurons and muscle. This gene encodes one member of the sodium channel alpha subunit gene family. It is heterogeneously expressed in the brain, and mutations in this gene have been linked to several

seizure disorders. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined.

To place an order, please Click HERE.

Catalog #:A10574

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:6326

Isotype:IgG

Swiss Prot:Q99250

Purity: Affinity purification

For research use only.





