

CCND1

Reactivity:Human Mouse Rat

Tested applications:WB IHC ICC IP

Recommended Dilution:WB 1:100 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IP 1:20 - 1:50

Calculated MW:34 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CCND1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

BCL1; PRAD1; U21B31; D11S287E;

Catalog #:A10757

Antibody Type:

Monoclonal Antibody

Species:Rabbit

Gene ID:595

Isotype:IgG

Swiss Prot:P24385

Purity:Affinity purification

For research use only.

Background:

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with tumor suppressor protein Rb and the expression of this gene is regulated positively by Rb. Mutations, amplification and overexpression of this gene, which alters cell cycle progression, are observed frequently in a variety of tumors and may contribute to tumorigenesis.

To place an order, please [Click HERE](#).