## CD4

Reactivity: Human

Tested applications: WB IHC ICC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:51 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CD4

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

CD4mut:

Monoclonal Antibody

Species:Rabbit

Gene ID:920

Isotype:IgG

Swiss Prot:P01730

Purity: Affinity purification

For research use only.

## Background:

This gene encodes a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigenes and is also a receptor for the human immunodeficiency virus. This gene is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified in this gene.

To place an order, please Click HERE.





