

MAPK14

Reactivity:Human Mouse Rat

Tested applications:WB ICC IF

Recommended Dilution:WB 1:500 - 1:2000 ICC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:38 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MAPK14

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

RK; p38; CSBP; EXIP; Mxi2; CSBP1; CSBP2; CSPB1; PRKM14; PRKM15; SAPK2A; p38ALPHA;

Background:

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various environmental stresses and proinflammatory cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its autophosphorylation triggered by the interaction of MAP3K7IP1/TAB1 protein with this kinase. The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress related transcription and cell cycle regulation, as well as in genotoxic stress response. Four alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

To place an order, please [Click HERE](#).

Catalog #:A10832

Antibody Type:

Monoclonal Antibody

Species:Rabbit

Gene ID:1432

Isotype:IgG

Swiss Prot:Q16539

Purity:Affinity purification

For research use only.