

MAGED1

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:86kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human MAGED1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

1mg/ml

Synonym:

MAGED1;DLXIN-1;NRAGE ;

Catalog #:A1092

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:9500

Isotype:IgG

Swiss Prot:Q9Y5V3

Purity:Affinity purification

For research use only.

Background:

MAGE-D1, also known as NRAGE or Dlxin-1, is a member of the MAGE family of proteins. Identified as a p75 neurotrophin receptor intracellular binding protein, MAGE-D1 induces developmental apoptosis of motoneurons, and is required for p75NTR-dependent apoptosis in sympathetic neurons. It is suggested that MAGE-D1 can suppress the motility and adhesion response of tumor cells. By forming a heterocomplex with Dlx/Msx family homeodomain proteins and Necdin, MAGE-D1 modulates the function of Dlx/Msx homeodomain during terminal differentiation and maturation of neurons. MAGE-D1 is also involved in the phosphorylation of IKK-alpha/beta, and subsequent transcriptional activation of the p65 subunit of NF-kappaB, via the XIAP-Tak1-Tab1 complex. (20100315, 19639218, 17453828, 15272023) G.

To place an order, please [Click HERE](#).