

## ATP2A2

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:** 115kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human ATP2A2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

i

**Synonym:**

ATP2A2; ATP2B; DAR; DD; DKFZp686P0211; FLJ20293; FLJ38063; MGC45367; SERCA2 ;

**Catalog #:** A1097

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 488

**Isotype:** IgG

**Swiss Prot:** P16615

**Purity:** Affinity purification

For research use only.

**Background:**

The ATP2A2 (SERCA2) calcium pump is one of several sarcoplasmic and endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPases responsible for regulating calcium transport across intracellular membranes (1). Multiple isoforms have been isolated, with ATP2A2a (SERCA2a) found predominantly in the sarcoplasmic reticulum of muscle cells and ATP2A2b (SERCA2b) more ubiquitously expressed in the endoplasmic reticulum of most cell types (2). An isoform containing a truncated carboxy region (ATP2A2c) is expressed in epithelial and hematopoietic cell lines and may be involved in monocyte differentiation (3). Post-translational modification of ATP2A2 (SERCA2), including phosphorylation and tyrosine nitration, modify  $\text{Ca}^{2+}$ -ATPase activity and calcium transport (4,5). Mutation in the corresponding ATP2A2 (SERCA2) gene results in Darier disease, a skin disorder characterized by the presence of dark, keratotic papules or rash found on the head and torso (6).

**To place an order, please [Click HERE](#).**