

Hydroxyl-Histone H2A-Y39

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:14 kDa

Observed MW:Refer to figures

Immunogen:

A specific peptide of Hydroxyl-Histone H2A-Y39

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

H2A.1; H2A.2; H2A/a; H2AFA;

Catalog #:A11009

Antibody Type:

Monoclonal Antibody

Species:Rabbit

Gene ID:3012

Isotype:IgG

Swiss Prot:P04908

Purity:Affinity purification

For research use only.

Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

To place an order, please [Click HERE](#).