MMP14

Reactivity: Human Mouse Rat

Tested applications: WB IHC ICC IF IP FC

Recommended Dilution: WB 1:500 - 1:1000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100

IP 1:20 - 1:50 FC 1:20 - 1:50

Calculated MW:65 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MMP14

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

MMP-14; MMP-X1; MT-MMP; MT1MMP; MTMMP1; WNCHRS; MT1-MMP; MT-MMP 1;

Background:

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the protein encoded by this gene is a member of the membrane-type MMP (MT-MMP) subfamily; each member of this subfamily contains a potential transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. This protein activates MMP2 protein, and this activity may be involved in tumor invasion.

To place an order, please Click HERE.

Monoclonal Antibody

Species: Rabbit

Gene ID:4323

Isotype:IgG

Swiss Prot:P50281

Purity: Affinity purification

For research use only.





