

## Tp53

**Reactivity:**Human

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

**Calculated MW:**44kDa

**Observed MW:**Refer to Figures

**Immunogen:**

A synthetic peptide of human Tp53

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

bb1; bfy; bhy; p44; p53; Tp53

**Catalog #:**A11232

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**7157

**Isotype:**IgG

**Swiss Prot:**P02340

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mice deficient for this gene are developmentally normal but are susceptible to spontaneous tumors. Evidence to date shows that this gene contains one promoter, in contrast to alternative promoters of the human gene, and transcribes a few of splice variants which encode different isoforms, although the biological validity or the full-length nature of some variants has not been determined.

**To place an order, please [Click HERE](#).**