www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

PIK3R3

Reactivity: Human Mouse Rat

Tested applications: WB IHC ICC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:54 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human PIK3R3

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

p55; p55PIK; p55-GAMMA;

Background:

Phosphatidylinositol 3-kinase (PI3K) phosphorylates phosphatidylinositol and similar compounds, which then serve as second messengers in growth signaling pathways. PI3K is composed of a catalytic and a regulatory subunit. The protein encoded by this gene represents a regulatory subunit of PI3K. The encoded protein contains two SH2 domains through which it binds activated protein tyrosine kinases to regulate their activity.

To place an order, please Click HERE.

Antibody Type:

Monoclonal Antibody

Species:Rabbit

Gene ID:8503

Isotype:IgG

Swiss Prot:Q92569

Purity: Affinity purification

For research use only.







