www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

MAP1LC3A

Reactivity: Human Mouse Rat

Tested applications: WB IHC ICC IF IP FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 IF 1:50 - 1:100

IP 1:20 - 1:50 FC 1:20 - 1:50

Calculated MW:14 kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human MAP1LC3A

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3;

Background:

MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. The protein encoded by this gene is one of the light chain subunits and can associate with either MAP1A or MAP1B. Two transcript variants encoding different isoforms have been found for this gene. The expression of variant 1 is suppressed in many tumor cell lines, suggesting that may be involved in carcinogenesis.

To place an order, please Click HERE.



Catalog #:A11428

Antibody Type:

Monoclonal Antibody

Species: Rabbit

Gene ID:84557

Isotype:IgG

Swiss Prot:Q9H492

Purity: Affinity purification

For research use only.





