

F8

Reactivity:Human Mouse Rat

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000

Calculated MW:267kDa/25kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human F8

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

AHF; F8B; F8C; HEMA; FVIII; DXS1253E;

Catalog #:A1366

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:2157

Isotype:IgG

Swiss Prot:P00451

Purity:Affinity purification

For research use only.

Background:

This gene encodes coagulation factor VIII, which participates in the intrinsic pathway of blood coagulation; factor VIII is a cofactor for factor IXa which, in the presence of Ca⁺² and phospholipids, converts factor X to the activated form Xa. This gene produces two alternatively spliced transcripts. Transcript variant 1 encodes a large glycoprotein, isoform a, which circulates in plasma and associates with von Willebrand factor in a noncovalent complex. This protein undergoes multiple cleavage events. Transcript variant 2 encodes a putative small protein, isoform b, which consists primarily of the phospholipid binding domain of factor VIIIc. This binding domain is essential for coagulant activity. Defects in this gene results in hemophilia A, a common recessive X-linked coagulation disorder. [provided by RefSeq, Jul 2008]

To place an order, please [Click HERE](#).