

KDR

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF IP

Recommended Dilution:WB 1:500 - 1:1000 IHC 1:50 - 1:500 IF 1:50- 1:500 IP 1:20- 1:100

Calculated MW:152kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant Protein of human KDR

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

a

Synonym:

FLK1; CD309; VEGFR; VEGFR2

Catalog #:A1484

Antibody Type:

Monoclonal Antibody

Species:Mouse

Gene ID:3791

Isotype:IgG

Swiss Prot:P35968

Purity:Affinity purification

For research use only.

Background:

KDR has also been designated as VEGFR-2 (Vascular endothelial growth factor receptor 2), CD309 (cluster of differentiation 309) and Flk1 (fetal liver kinase 1). Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. KDR is one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.

To place an order, please [Click HERE](#).