

## MMP13

**Reactivity:** Human Mouse

**Tested applications:** WB

**Recommended Dilution:** WB 1:500 - 1:2000

**Calculated MW:** 54kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human MMP13

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CLG3; MANDP1;

**Catalog #:** A1606

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 4322

**Isotype:** IgG

**Swiss Prot:** P45452

**Purity:** Affinity purification

For research use only.

**Background:**

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The protein encoded by this gene cleaves type II collagen more efficiently than types I and III. It may be involved in articular cartilage turnover and cartilage pathophysiology associated with osteoarthritis. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq, Jul 2008]

**To place an order, please [Click HERE](#).**