

BTRC

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

Calculated MW:69kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human BTRC

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

dmpt

Synonym:

BETA-TRCP; FBW1A; FBXW1; FBXW1A; FWD1; MGC4643; bTrCP; bTrCP1; betaTrCP;

Catalog #:A1656

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8945

Isotype:IgG

Swiss Prot:Q9Y297

Purity:Affinity purification

For research use only.

Background:

-transducin repeat-containing protein (-TrCP or FBW1A) is an F-box family protein characterized by the presence of the protein-protein mediating F-box domain first described in cyclin F. F-box proteins act as substrate adaptors that target proteins containing a specific phosphorylated sequence element, referred to as a phosphodegron, to the SCF E3 ubiquitin ligase complex for ubiquitination (1,2). -TrCP targets many important proteins with diverse functions, such as p53, H-Ras, Smad4, IB, -catenin, and the cell cycle checkpoint protein claspin, for ubiquitin-mediated degradation (3-5). Research studies have shown that inhibition of -TrCP expression has a demonstrated benefit in the treatment of prostate cancer (6).

To place an order, please [Click HERE](#).