

EWSR1

Reactivity:Human Mouse

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:68kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human EWSR1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

EWS; bK984G1.4;

Catalog #:A1690

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:2130

Isotype:IgG

Swiss Prot:Q01844

Purity:Affinity purification

For research use only.

Background:

This gene encodes a multifunctional protein that is involved in various cellular processes, including gene expression, cell signaling, and RNA processing and transport. The protein includes an N-terminal transcriptional activation domain and a C-terminal RNA-binding domain. Chromosomal translocations between this gene and various genes encoding transcription factors result in the production of chimeric proteins that are involved in tumorigenesis. These chimeric proteins usually consist of the N-terminal transcriptional activation domain of this protein fused to the C-terminal DNA-binding domain of the transcription factor protein. Mutations in this gene, specifically a t(11;22)(q24;q12) translocation, are known to cause Ewing sarcoma as well as neuroectodermal and various other tumors. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1 and 14.

To place an order, please [Click HERE](#).