

RBX1

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:12kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human RBX1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

1mg/ml

Synonym:

BA554C12.1; MGC13357; MGC1481; RNF75; ROC1;

Catalog #:A1832

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:9978

Isotype:IgG

Swiss Prot:P62877

Purity:Affinity purification

For research use only.

Background:

RING-box protein 1 (RBX1 or ROC1) is an essential component of two distinct but structurally related E3 ubiquitin ligase complexes, the SCF complex and the CBC (VHL) complex (1). RBX1 mediates the neddylation of CUL1, which activates SCF E3 ligase by facilitating the ubiquitin transfer from E2 to substrates (2-4). The RING finger domain of RBX1 is required for ubiquitin ligation (5). Two evolutionarily conserved mammalian RBX family members, RBX1/ROC1 and RBX2/ROC2/SAG, have been identified (5). RBX1 is constitutively expressed and binds to CUL2/VHL, while stress-inducible RBX2 binds to CUL5/SOCS (6).

To place an order, please [Click HERE](#).