

AREG

Reactivity: Human Mouse

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 28kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human AREG

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

AR; CRDGF; MGC13647; SDGF;

Catalog #: A1860

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 374

Isotype: IgG

Swiss Prot: P15514

Purity: Affinity purification

For research use only.

Background:

Amphiregulin (AREG) is one of the ligands of the epidermal growth factor receptor (EGFR). AREG plays a central role in mammary gland development and branching morphogenesis in organs and is expressed both in physiological and in cancerous tissues. The AREG protein is synthesized as a 252-amino acid transmembrane precursor, pro-AREG. At the plasma membrane, pro-AREG is subjected to sequential proteolytic cleavages within its ectodomain and is then released as the soluble AREG protein. Depending on the cell type and microenvironment, AREG can be produced in multiple cellular and mature forms using alternative pro-AREG cleavage sites and glycosylation motifs.

To place an order, please [Click HERE](#).