

PDHA1

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:43kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human PDHA1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

PDHA; PDHCE1A; PHE1A;

Catalog #:A1895

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:5160

Isotype:IgG

Swiss Prot:P08559

Purity:Affinity purification

For research use only.

Background:

The pyruvate dehydrogenase complex catalyzes the conversion of pyruvate and CoA into acetyl-CoA and CO₂ in the presence of NAD⁺. Acetyl-CoA then goes into the citric acid cycle where it reacts with oxaloacetate to form citrate. Acetyl-CoA is also used for fatty acid and cholesterol biosynthesis. The reaction of oxidative decarboxylation of pyruvate therefore serves as a critical link between glycolysis and the citric acid cycle and lipid metabolism. In mammalian cells, the pyruvate dehydrogenase complex is located in the mitochondrial matrix (1). This complex is comprised of three enzymes: pyruvate dehydrogenase (E1), dihydrolipoamide acetyltransferase (E2) and dihydrolipoamide dehydrogenase (E3). Pyruvate dehydrogenase (E1) consists of two subunits: and . This enzyme catalyzes the removal of CO₂ from pyruvate. Mutations in the subunits of pyruvate dehydrogenase (E1) lead to congenital defects that are usually associated with lactic acidosis, neurodegeneration and early death (2).

To place an order, please [Click HERE](#).