

CCL21

Reactivity: Human

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW: 15kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human CCL21

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

ECL; SLC; CKb9; TCA4; 6Ckine; SCYA21;

Catalog #: A1896

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 6366

Isotype: IgG

Swiss Prot: O00585

Purity: Affinity purification

For research use only.

Background:

This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. Similar to other chemokines the protein encoded by this gene inhibits hemopoiesis and stimulates chemotaxis. This protein is chemotactic in vitro for thymocytes and activated T cells, but not for B cells, macrophages, or neutrophils. The cytokine encoded by this gene may also play a role in mediating homing of lymphocytes to secondary lymphoid organs. It is a high affinity functional ligand for chemokine receptor 7 that is expressed on T and B lymphocytes and a known receptor for another member of the cytokine family (small inducible cytokine A19).

To place an order, please [Click HERE](#).