

HIST2H2BE

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:14kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human HIST2H2BE

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

H2B; H2BQ; GL105; H2B.1; H2BFQ; H2BGL105;

Catalog #:A1958

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8349

Isotype:IgG

Swiss Prot:Q16778

Purity:Affinity purification

For research use only.

Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a member of the histone H2B family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

To place an order, please [Click HERE](#).