PCK1

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:1000 - 1:2000 IHC 1:50 - 1:200

Calculated MW:69kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human PCK1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonym:

MGC22652; PEPCK-C; PEPCK1; PEPCKC;

Gene ID:5105

Isotype:IgG

Swiss Prot:P35558

Purity: Affinity purification

For research use only.

Background:

PCK1(Phosphoenolpyruvate carboxykinase, cytosolic) is also named as PEPCK1 and belongs to the phosphoenolpyruvate carboxykinase [GTP] family. It catalyzes the formation of phosphoenolpyruvate from oxaloacetate, with the release of carbon dioxide and GDP. It is also a main control point for the regulation of gluconeogenesis. In eukaryotes there are two isozymes: a cytoplasmic one and a mitochondrial one. Defects in PCK1 are the cause of cytosolic phosphoenolpyruvate carboxykinase deficiency (C-PEPCKD). This antibody is specific to PCK1.

To place an order, please Click HERE.



