

## ICAM2

---

**Reactivity:**Human

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**31kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human ICAM2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

CD102; ICAM2;

**Catalog #:**A2038

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**3384

**Isotype:**IgG

**Swiss Prot:**P13598

**Purity:**Affinity purification

For research use only.

**Background:**

The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM) family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9 immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell adhesion. It mediates adhesive interactions important for antigen-specific immune response, NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions important for immune response and surveillance. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2008]

**To place an order, please [Click HERE](#).**