

SLAMF1

Reactivity: Human

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 70kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human SLAMF1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CD150; CDw150; SLAM;

Catalog #: A2044

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 6504

Isotype: IgG

Swiss Prot: Q13291

Purity: Affinity purification

For research use only.

Background:

Following occupancy of the T cell receptor by antigen, T cell proliferation and lymphokine production are determined by a second costimulatory signal delivered by a ligand expressed on antigen-presenting cells. SLAM (for signaling lymphocyte-activation molecule, also designated CDw150) is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. SAP (for SLAM-associated protein) contains an SH2 domain and functions to inhibit SH-PTP2 recruitment to the SLAM docking site, an activity induced by Fyn phosphorylation of SLAM. Mutations of the SAP gene may be associated with X-linked lympho-proliferative disease (XLP).

To place an order, please [Click HERE](#).